


Background

- Coronavirus of 2019 (COVID-19) is an infectious virus currently causing a pandemic. Evidence supports high rates of asymptomatic infections with COVID-19, leading to concerns that viral transmission may occur via these asymptomatic patients.
- Initially, the screening of asymptomatic patients was of low utility as only 2 of 1,342 ED patients tested positive. These findings were during a period of low COVID-19 prevalence in the community.
- Subsequently, COVID-19 infection rates increased in the community. During this increase in COVID-19, the rate of asymptomatic cases also increased.



Aims

To describe the utility of screening in the Emergency Department for COVID-19 in asymptomatic patients.

Methods

- Retrospective cohort study that included all adult ED patients who had a COVID-19 test via the ED's asymptomatic screening protocol. The dates of data collected ranged from the May 1st, 2020 to October 31st, 2020.
- The primary outcome was positive COVID-19 tests. Patients with positive tests underwent further data abstraction using standard chart review methodology. Data were described with simple descriptive statistics.
- Signs/symptoms of COVID-19 : fever, hypoxia, cough, fatigue, myalgias, nasal congestion/drainage, shortness of breath, sore throat, headache, diarrhea, or loss of taste/smell.
- Other data collected included: COVID risk factors, laboratory testing including COVID antibody, imaging findings if obtained, infectious disease consult, and treatments.

Results

Patients

- Total of 7408 adult patients had COVID test under the asymptomatic protocol
 - Mean age 53.8 ± 19.3 years
- Total of 78 positive tests (1.1%, 95%CI 0.8-1.3) and 9 inconclusive.
- 48 (62%) underwent infectious disease consult.
- No patient died due to COVID.




Figure 1. Of the positive cases, 32 (41%, 95%CI 30-53) had signs/symptoms (n=29) or were unable to be assessed due to decreased mental status (n=3).

Figure 2. 13 patients received specific therapy following positive COVID test

Therapy	Supplemental oxygen	Dexamethasone	Convalescent plasma	Remdesivir
# of patients	12	6	6	4

Conclusion

Asymptomatic COVID-19 testing is of low yield when the daily community rate is less than 4 cases/100,000 individuals. Of those found to be positive under an asymptomatic testing protocol, nearly half had signs or symptoms attributed to COVID-19 infection suggesting the rate of asymptomatic infection was lower than reported. In the current pandemic, ED physicians should carefully assess for COVID-19 signs and symptoms even when patients present for other indications.

Acknowledgments

Thank you to Dr. James Holmes for the mentorship for my Scholarly Project Option (SPO), creation and submission of the abstract, and guidance throughout.

Questions or comments?
Please contact: Evan Chua, MS4
Email: ecchua@ucdavis.edu

